

101587,372

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
13 June 2002 (13.06.2002)

PCT

(10) International Publication Number
WO 02/45485 A1

- (51) International Patent Classification⁷: A01H 5/00, Rosella Street, Nicholls, Australian Capital Territory 2913 (AU).
C12N 15/11, 15/29, 15/54
- (21) International Application Number: PCT/AU01/01580 (74) Agents: OLIVE, Mark, R. et al.; FB RICE & CO, 139 Rathdowne Street, Carlton, Victoria 3053 (AU).
- (22) International Filing Date: 7 December 2001 (07.12.2001) (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (25) Filing Language: English (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (26) Publication Language: English
- (30) Priority Data:
PR 1975 8 December 2000 (08.12.2000) AU
60/251,852 8 December 2000 (08.12.2000) US
- (71) Applicant (for all designated States except US): COMMONWEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANISATION [AU/AU]; Limestone Avenue, Campbell, Australian Capital Territory 2601 (AU).
- (72) Inventors; and
(75) Inventors/Applicants (for US only): LLWELLYN, Danny [AU/AU]; 6 Myall Street, O'Connor, Australian Capital Territory 2602 (AU). FURBANK, Robert [AU/AU]; 35 Gillespie Street, Weetangara, Australian Capital Territory 2614 (AU). RUAN, Yong-Ling [AU/AU]; 9
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/45485 A1

(54) Title: MODIFICATION OF SUCROSE SYNTHASE GENE EXPRESSION IN PLANT TISSUE AND USES THEREFOR

(57) Abstract: Methods and means are provided to modulate fibre quality in fibre-producing plants, such as cotton by modulating sucrose synthase activity and/or expression in such plants. The methods and means may also be used to obtain plants with seedless fruits or male-sterile plants.

Modification of sucrose synthase gene expression in plant tissue and uses therefor

Technical Field

The present invention relates to modifying targeted gene expression in plants to obtain a desired effect in the plant.

5

Background Art

Sucrose synthase (SuSy) is a key enzyme in the breakdown of sucrose in all plant sink tissue, including grain and fruit and has been extensively studied in many plants. Only relatively recently, however, has this protein and gene been characterised from cotton. The full length (2625 bp) of cotton sucrose synthase (SuSy) was isolated by Perez-Grau, L and Delmer, D in UC, Davis (accession number U73588) in May 1996. A 2030 bp fragment of the same cDNA with 595 bp missing at the 5' end was isolated by the same group in 1994 and was given to Prem Chourey in USDA/ARS for collaborative research. However, no evidence was available at that time regarding the role of this SuSy gene in cotton fibre/seed development, although it had been speculated (Amor *et al.*, 1995) that part of the fibre localised SuSy could associate with cellulose synthase playing a role to channel carbon to this enzyme.

20 Evidence has been obtained that the expression of the SuSy gene could be important not only for cellulose synthesis but also for fibre cell initiation (thus may control fuzz) and a model on how sucrose is partitioned and competed for between fibre, seed coat and embryos of the cotton seed was proposed (Ruan *et al* 1997 *Plant Physiol.* 115, 375-385; Ruan and Chourey 1998 *Plant Physiol.* 25 118, 399-406). More recently, the present inventors obtained further evidence supporting the hypothesis that SuSy plays a key role in mobilising sucrose into initiating fibre cells (Ruan *et al.*, 2000 *Aust. J. Plant Physiol.* 27, 795-800).

The art is thus deficient in providing methods and means for altering the fibre development and properties in plants, particularly cotton, through alteration of

30

sucrose synthase levels in cells of the plants. The present inventors have now shown that it is possible to modify one or more fibre characteristics and/or fibre content by modifying SuSy gene expression, particularly by modifying SuSy expression in transgenic cotton.

5

These and other problems are solved as described hereinafter in the different embodiments and claims.

Throughout this specification, unless the context requires otherwise the word
10 "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps but not the exclusion of any other element, integer or step or group of elements, integers or steps.

15 Summary of the invention.

The invention provides a method for altering fibre development or properties of a fibre producing plant, preferably a cotton plant, such as, for example, a cotton plant of the variety fribremax and particularly the Fibermax™ variety, said method comprising the steps of providing cells of said plants with a chimeric
20 gene comprising the following operably linked DNA fragments: a plant operable promoter, preferably a subterranean clover stunt virus promoter; a coding region which when transcribed yields an RNA said RNA being capable of reducing the expression of an endogenous sucrose synthase gene, preferably an endogenous sucrose synthase gene expressed in fibre cells, preferably fibre
25 initial cells, or capable of being translated into an active sucrose synthase protein; and transcription termination and polyadenylation signals that function in the plant cells.

In the embodiments where the RNA is capable of being translated into an
30 active sucrose synthase protein preferred coding regions comprise a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2;

(b) a nucleotide sequence comprising the nucleotide sequence of SEQ ID NO: 1; (c) a nucleotide sequence having at least 70% sequence identity with the nucleotide sequence of (a) or (b); and (d) a nucleotide sequence hybridizing under stringent conditions with the complement of (a) or (b); and (e) a fragment
5 of any one of (a) through (d) encoding an active sucrose synthase.

In the embodiments where the RNA is capable of reducing the expression of an endogenous sucrose synthase gene, preferred coding regions comprise a nucleotide sequence selected from the group consisting of: (a) a nucleotide
10 sequence comprising at least about 19 or 25 contiguous nucleotides having at least 70% sequence identity to a nucleotide sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2 or the complement thereof; and (b) a nucleotide sequence comprising at least about 19 or 25 contiguous nucleotides having at least 70% sequence identity from a nucleotide
15 sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 1 or the complement thereof. More particularly, the coding region comprises the nucleotide sequence of SEQ ID NO: 1 from the nucleotide at about position 2208 to the nucleotide at about position 2598, or the complement thereof. The coding region may comprise simultaneously both
20 sense and antisense nucleotide sequences capable of forming a double stranded RNA molecule.

It is another objective of the invention to provide a method for improving fibre yield in a fibre producing plant, comprising the steps of providing cells of said
25 plant with a chimeric gene comprising the following operably linked DNA fragments: (a) a plant operable promoter; (b) a DNA region capable of being translated into an active sucrose synthase protein; and (c) transcription termination and polyadenylation signals that function in said plant cells.

30 The invention also provides a method for improving fibre quality in a fibre producing plant, comprising the steps of providing cells of said plant with a

chimeric gene comprising the following operably linked DNA fragments: (a) a plant operable promoter; (b) a DNA region capable of being translated into an active sucrose synthase protein; and (c) transcription termination and polyadenylation signals that function in said plant cells.

5

In yet another embodiment of the invention a method is provided for increasing seed size in a fibre producing plant, comprising the steps of providing cells of said plant with a chimeric gene comprising the following operably linked DNA fragments: (a) a seed-specific promoter; (b) a DNA region capable of being translated into an active sucrose synthase protein; and (c) transcription termination and polyadenylation signals that function in said plant cells.

10

The invention also relates to fibre producing plants, particularly cotton plants, comprising in their genome a chimeric DNA comprising the following operably linked DNA fragments: (a) a plant operable promoter; (b) a coding region which when transcribed yields an RNA, said RNA being capable of reducing the expression of an endogenous sucrose synthase gene, preferably an endogenous sucrose synthase gene expressed in fibre cells, preferably fibre initial cells, or alternatively, which is capable of being translated into an active sucrose synthase protein; and (c) transcription termination and polyadenylation signals that function in said plant cells.

15

20

Yet another objective of the invention is to provide a fibre producing plant comprising a chimeric DNA according to the invention wherein said DNA is transcribed to yield RNA that is capable of increasing the expression of an endogenous sucrose synthase gene, preferably an endogenous sucrose synthase gene expressed in fibre cells, preferably fibre initial cells and said fibre cells have an increased sucrose synthase activity compared to fibre cells of plant cells which do not comprise said chimeric DNA.

25

30

Yet another objective of the invention is to provide a fibre producing plant comprising a chimeric DNA according to the invention, wherein said chimeric DNA is transcribed to yield RNA that is capable of being translated into an active sucrose synthase protein, thereby increasing the level of sucrose synthase in the plant relative to a plant that does not comprise said chimeric DNA. Preferably, the active sucrose synthase is expressed in fibre cells, more preferably fibre initial cells, wherein said fibre cells preferably have higher sucrose synthase activity compared to the fibre cells of a plant which does not comprise the chimeric DNA.

10

In yet another embodiment of the invention fibre producing plants are provided comprising a chimeric DNA according to the invention, wherein said coding region comprises a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2; (b) a nucleotide sequence comprising the nucleotide sequence of SEQ ID NO: 1; (c) a nucleotide sequence having at least 70% sequence identity with the nucleotide sequence of (a) or (b); (d) a nucleotide sequence hybridizing under stringent conditions with the nucleotide sequence of (a) or (b); and (e) a fragment of any one of (a) through (d) encoding an active sucrose synthase.

20

The invention also provides a fibre producing plant comprising the chimeric DNA of the invention wherein said chimeric DNA encodes an RNA capable of reducing an endogenous sucrose synthase gene expression, such that said fibre cells have a reduced sucrose synthase activity compared to fibre cells of plant cells that do not comprise said chimeric DNA, particularly wherein the coding region comprises a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence comprising at least about 19 or 25 contiguous nucleotides having at least 70% sequence identity to a nucleotide sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2, or the complement thereof; and (b) a nucleotide sequence

25

30

comprising at least about 19 or 25 contiguous nucleotides having at least 70% sequence identity from a nucleotide sequence encoding a polypeptide comprising the nucleotide sequence of SEQ ID NO: 1 or the complement thereof; and (c) the nucleotide sequence of SEQ ID NO: 1 from the nucleotide
5 at about position 2208 to the nucleotide at about position 2598 or the complement thereof.

Also provided are seeds of a fibre producing plant, preferably the seed of a cotton plant comprising the chimeric DNA or genes according the invention, as
10 well as fibres with altered development or properties, isolated from such plants.

The invention also relates to the use of a sucrose synthase encoding nucleotide sequence for altering fibre yield or fibre quality or fibre properties.

15 Brief Description of Drawings

Figure 1 is a photographic representation showing that down regulation of SuSy in the fibre causes a decrease in fibre elongation in transgenic plants. Transverse sections of cotton bolls and seeds of wild type (WT) and transgenic cotton carrying an antisense SuSy gene construct (SuSy/A) are shown. Three
20 developmental stages are shown: 0, 2 and 6 days after anthesis. Data indicate reduced fibre in transgenic lines, particularly in samples 2 days post anthesis (DAA) or older.

Figure 2 is a photographic representation of an immunohistochemical stain
25 showing localisation of SuSy protein in wild type (WT) cotton (panel A), and the transgenic lines 291-147 (panel B) and 295-82 (panel C). Black coloration in developing fibre cells on the surface of the ovule, and in the ovule epidermis, of the wild type (WT) plants indicates SuSy protein. Transgenic lines show reduced amount of SuSy protein (reduced black coloration). No SuSy protein
30 was detectable in line 295-82. This reduced SuSy protein correlates with lack of fibre development in the transgenic lines.

Figure 3 is a photographic representation of cotton bolls (left) and seed (right) from wild type (WT) cotton and the transgenic line 294-147 carrying an antisense SuSy gene construct, which has reduced SuSy protein in the embryo (upper boll and seeds). Seeds from the transgenic line are fibreless, but there is also a great reduction in seed size in seeds labeled III. Some seeds did not form at all.

Figure 4 is a photographic representation of wild type cotton plants (left of the figure) and two transformed cotton lines (plants at the middle and right of the figure). Data show that there was no detectable phenotypic effect of the transgene with respect to vegetative growth of plants.

Figure 5 is a schematic representation of an antisense gene construct (top) and a cosuppression gene construct (below) used to suppress SuSy gene expression *in planta*. Gene construct components are as follows: nptII, neomycin phosphotransferase marker gene for selection of transformants; S1 and S7, subclover stunt virus promoters 1 and 7, respectively (see Australian Patent No 689311); S3 and S5, subclover stunt virus gene terminator sequences 3 and 5, respectively (see Australian Patent No 689311); SuSy, a 390 bp nucleotide sequence of a sucrose synthase-encoding cDNA (Genbank Accession number U73588) from CTGGGAT to TGA CTT (ie. 211 bp coding region upstream of the stop codon plus 179 bp 3'-untranslated (UTR) region; nucleotide position 2208 to nucleotide position 2598 of SEQ ID NO: 1). Arrows indicate the orientation of fragments relative to their native orientation in the genes from which they were derived.

Figure 6 is a copy of a photographic representation of a Southern blot hybridization, showing SuSy gene copy in wild type cotton plants and transgenic cotton lines. DNAs were isolated from T0 generation plants (panel A), and a segregating population of T1 generation plants from the primary

transformed lines designated 14 and 4 (panel B). The arrows indicate lines 14, 8, and 4 in the T0 generation having a fibre-less or reduced fibre phenotype (panel A), and lines 14-1, 14-2, 14-3, 4-1, and 4-3 in the segregating population of T1 plants having a fibre-less or reduced fibre phenotype (panel B). The numbers indicate the transgenic cotton line from which the individual plants were derived.

Figure 7 is a photographic representation of a scanning electron microscopy (SEM) profile of the ovule epidermis of wild type cotton (panel A) and line 294-147 shown in Fig. 2B (panels B and C). Many fibre cells are collapsed and/or shrunk in the transgenic line, compared to the wild type plants. Whilst some fibre cells appeared in the transgenic lines, these were much smaller in size and fewer in number than in the wild type ovule (compare panels A and B). The shrunk and/or collapsed phenotype of the transgenic line was clearly demonstrated under high magnification (panel C).

Figures 8, 9 and 10 shows the correlation between changes of SuSy activity (Fig. 8) and reduction of fibre length (Fig. 9) in 3 day old seeds from 10 segregating individuals of three T1 generation lines (lines 147, 43 and 101), together with line 82 and wild type (WT). Figure 10 represents the regression analysis between SuSy activity and fibre length.

Figure 8 is a graphical representation showing modified SuSy activity (abscissa) in transgenic lines compared to wild type plants. Plant lines are indicated on the x-axis. Error bars represent the SEM for assays of 3-day old seeds from 10 segregating individuals of the T1 generation lines.

Figure 9 is a graphical representation showing modified fibre length (abscissa) in transgenic lines compared to wild type plants. Plant lines are indicated on the x-axis. Error bars represent the SEM for assays of 3-day old seeds from 10 segregating individuals of the T1 generation lines.

Figure 10 is a graphical representation of a regression analysis between SuSy activity data presented in Figure 8 (x-axis) and fibre length data presented in Figure 9 (abscissa). Data show a strong correlation ($R^2=0.98$) between SuSy activity and fibre length over the range shown.

Figure 11 is a photographic representation showing fuzz fibre length in seeds of wild type cotton (panel A) and the transgenic line 147#2 (panel B), and showing reduced fuzz in the transgenic line. Data demonstrate that a reduction in SuSy gene expression reduces fuzz. Seed were delinted by hand. The oval represents the chalazal end of the mature seeds. Seed from the transgenic line are more brown in appearance than wild type seeds, because the underlying brown seed coat is covered by shorter fuzz fiber than in wild type seed.

Detailed Description of the Invention

In a first aspect, the present invention provides a plant having altered expression of an iso-form of sucrose synthase (SuSy) resulting in an altered fibre, fruit or seed production ability.

Preferably, the plant is a transgenic plant having an under- or overexpression of an iso-form of SuSy involved in fruit/seed production. In a preferred form, the plant retains expression of SuSy in developing vegetative tissue such that growth and development of the plant is not adversely affected.

When the expression of an isoform of SuSy that is involved in fruit/seed production is reduced in the plant, preferably in the seed, more preferably in the fibre producing cells, even more preferably in fibre initial cells, the plant has a reduced fibre and/or seed production. When the expression of an isoform of SuSy that is involved in fruit/seed production is enhanced (i.e. SuSy is over-expressed) in the plant, preferably in the seed, more preferably in the fibre

cells, even more preferably in fibre initial cells, the plant has enhanced fibre and/or seed production.

The ability to cause underexpression of an iso-form of SuSy involved in fruit/seed production can be used to develop plants which produce few or no seeds but are still able to grow and produce fruit normally. The ability to cause overexpression of an iso-form of SuSy involved in fruit/seed production in a tissue specific manner can be used to develop plants which produce greater amounts of fibre or longer fibre or having altered fibre structure but are still able to grow and develop normally. The plant can be any plant in which the expression of an iso-form of SuSy is involved in fruit/seed production. The present invention is applicable for modifying a wide range of horticultural crops such as grape, peach, pear, and apple. More preferably, the plant is a cotton plant.

15

In a preferred embodiment, plants, particularly cotton plants, are provided comprising a chimeric gene, preferably stably integrated in their genome, the chimeric gene comprising the following operably linked DNA fragments:

- a) a plant operable promoter
- 20 b) a coding region which when transcribed yields an RNA which is either
 - i) capable of reducing the expression of an endogenous sucrose synthase gene, preferably an endogenous sucrose synthase gene expressed in fibre initial cells; or
 - ii) capable of being translated into an active sucrose synthase
- 25 c) transcription termination and polyadenylation signals that function in plant cells.

A particularly preferred embodiment of the coding region as defined sub ii) is a coding region comprising a nucleotide sequence which encodes a protein comprising the amino acid of SEQ ID NO: 2, particularly the nucleotide

30

sequence of SEQ ID NO: 1, or a fragment of such a nucleotide sequence capable of being translated into a functional sucrose synthase. The coding region encoding the sucrose synthase activity may be derived from other plant species or other species (as indicated elsewhere in this application), such as, 5 for example, from potato (cDNA sequence is available from Genbank library, accession number M18745).

A particularly preferred embodiment of the coding region as defined sub i) is a coding region comprising the complement of a nucleotide sequence 10 ("antisense") encoding a polypeptide encoded by the nucleotide sequence of SEQ ID NO: 1 from the nucleotide at about position 2208 to the nucleotide at about position 2598, particularly the coding region comprises the nucleotide sequence of SEQ ID NO: 1 from the nucleotide at position 2208 to the nucleotide at position 2598.

15

As used herein, the term "coding region" shall be taken to mean a DNA sequence which is capable of being transcribed into a biologically active RNA, irrespective of whether or not said DNA sequence encodes an amino acid sequence. In the present context, a biologically active RNA includes RNA 20 which is capable of inducing a biological effect in the target cell, such as antisense or sense RNA capable of triggering post transcriptional gene silencing, or a ribozyme etc.

Biologically active RNA also includes RNA which is capable of being translated 25 into a polypeptide or protein.

Having read these preferred embodiments, the person skilled in the art will immediately realize that functional equivalents of the mentioned coding regions may be used to similar effect.

30

Functional equivalents of a coding region capable of being transcribed into RNA which is capable of reducing the expression of an endogenous sucrose synthase gene, include e.g. shorter antisense fragments compared to the above-mentioned nucleotide sequences.

5

The length of the antisense nucleotide sequence may vary from about 10 nucleotides (nt) or 19 nt, up to a length equaling the length (in nucleotides) of the target nucleic acid (i.e. the full-length sucrose synthase gene). Preferably, the total length of the antisense nucleotide sequence is at least 10 nt, 10 preferably 15 nt or 19 nt, more preferably at least about 50 nt, more preferably at least about 100 nt, more preferably at least about 150 nt, more preferably at least about 200 nt, and even more preferably at least about 500 nt. It is expected that there is no upper limit to the total length of the antisense nucleotide sequence, other than the total length of the target nucleic acid. 15 However for practical reasons (such as e.g. stability of the chimeric genes) it is expected that the length of the antisense nucleotide sequence should not exceed 5000 nt, and preferably should not exceed 2500 nt. Conveniently, the length of an antisense nucleic acid is limited to about 1000 nt.

20 It will be appreciated that the longer the total length of the antisense nucleotide sequence is, the less stringent the requirements for sequence identity between the total antisense nucleotide sequence and the complement of the corresponding sequence in the target sucrose synthase gene become.

25 Preferably, the total antisense nucleotide sequence should have a sequence identity of at least about 75% with the complement corresponding target sequence, particularly at least about 80 %, more particularly at least about 85%, quite particularly about 90%, especially about 95%, more especially about 100%, quite especially be identical to complement of the corresponding part of 30 the target nucleic acid. However, it is preferred that the antisense nucleotide sequence always includes a sequence of about 10 consecutive nucleotides,

particularly about 19 or 20 nt, more particularly about 50 nt, especially about 100 nt, quite especially about 150 nt with 100% sequence identity to the complement of the corresponding part of the target nucleic acid. Preferably, for calculating the sequence identity and designing the corresponding antisense sequence, the number of gaps should be minimized, particularly for the shorter antisense sequences.

Particularly preferred are antisense nucleotide sequences having a sequence identity of at least about 75%, preferably at least about 80%; particularly at least about 85%; quite particularly about 90%, especially about 95% with the complement of the corresponding part of the nucleotide sequence of SEQ ID NO: 1.

For the purpose of this invention, the "sequence identity" of two related nucleotide or amino acid sequences, expressed as a percentage, refers to the number of positions in the two optimally aligned sequences which have identical residues (x100) divided by the number of positions compared. A gap, i.e. a position in an alignment where a residue is present in one sequence but not in the other is regarded as a position with non-identical residues. The alignment of the two sequences is performed by the Needleman and Wunsch algorithm (Needleman and Wunsch 1970) Computer-assisted sequence alignment, can be conveniently performed using standard software program such as GAP which is part of the Wisconsin Package Version 10.1 (Genetics Computer Group, Madison, Wisconsin, USA) using the default scoring matrix with a gap creation penalty of 50 and a gap extension penalty of 3.

It goes without saying that sense fragments may also be used for reducing the expression of an endogenous sucrose synthase gene, and the same embodiments of length and sequence homology as herein described for antisense molecules, apply mutatis mutandis to the sense molecules.

Particularly suited for reducing the expression of an endogenous sucrose synthase genes are DNA regions, preferably under the control of a plant-operable promoter, which when transcribed result in so-called double stranded RNA molecules, comprising both sense and antisense sequences which are capable of forming a double stranded RNA molecule as described in WO 99/53050 (incorporated herein in its entirety by reference). In this particular case, a chimeric gene may thus be introduced into a plant cell comprising a plant operable promoter operably linked to a DNA region, whereby that DNA region comprises a part of coding region comprising at least 10 or 19 consecutive nucleotides from the coding region of a nucleic acid encoding a sucrose synthase protein, such as but not limited to, a sucrose synthase protein with the amino acid sequence of SEQ ID NO: 1 (the so-called sense part) as well as a DNA sequence which comprises at least the complementary DNA sequence of at least 10 or 19 nucleotides of the sense part, but which may be completely complementary to the sense part (the so-called antisense part). The chimeric gene may comprise additional regions, such as a transcription termination and polyadenylation region functional in plants. When transcribed an RNA can be produced which may form a double stranded RNA stem between the complementary parts of the sense and antisense region. A spacer region may be present between the sense and antisense nucleotide sequence. The chimeric gene may further comprise an intron sequence, preferably located in the spacer region.

Functional equivalents of coding regions capable of being transcribed into RNA which can be translated into an active sucrose synthase comprise mutant or allelic forms derived from sucrose synthase genes, particularly from sucrose synthase gene with active expression in fibre initials.

Methods to derive mutants e.g. of a sucrose synthase gene, particularly of sucrose synthase genes encoding a protein as represented in SEQ ID NO: 2, quite particularly of sucrose synthase gene comprising the nucleotide sequence

of SEQ ID NO: 1, such a site-specific mutagenesis methods are well known in the art, as well as assays to identify active sucrose synthase enzymes encoded by the mutant sequences.

5 Allelic variants of the nucleotide sequences encoding sucrose synthase may be identified by hybridization of libraries, under stringent conditions, such as cDNA or genomic libraries of a different varieties or plant lines, particularly cotton varieties and plant lines. Nucleotide sequences which hybridize under stringent conditions to nucleotide sequences encoding the amino acid sequence of SEQ
10 ID NO: 2 or to the nucleotide sequence of SEQ ID NO: 1, or a sufficiently large part thereof (preferably about 25 contiguous nucleotides, particularly at least about 50 contiguous nucleotides, more particularly at least about 100 contiguous nucleotides) and which encode a functional protein with sucrose synthase activity are functional equivalents of the above mentioned preferred
15 coding regions. Such nucleotides may also be identified and isolated using e.g. polymerase chain reaction amplification using an appropriate pair of oligonucleotides having at least about 25 contiguous nucleotides, particularly at least about 50 contiguous nucleotides, more particularly at least about 100 contiguous nucleotides of the nucleotide of SEQ ID NO: 1.

20 "Stringent hybridization conditions" as used herein mean that hybridization will generally occur if there is at least 95% and preferably at least 97% sequence identity between the probe and the target sequence. Examples of stringent hybridization conditions are overnight incubation in a solution comprising 50%
25 formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared carrier DNA such as salmon sperm DNA, followed by washing the hybridization support in 0.1 x SSC at approximately 65 °C. Other hybridization and wash conditions are well known and are exemplified in
30 Sambrook *et al*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY (1989), particularly chapter 11.

The following are accession numbers of nucleotide sequences in the Genbank library which are plant sucrose synthase genes, parts thereof or nucleotide sequences having sequence similarity to sucrose synthase genes which could be used according to the methods herein described: BM094593 (Glycine max), BM093753 (Glycine max) BM093158 (Glycine max) BM092695(Glycine max) BM092443(Glycine max) BM092322(Glycine max) BM085310(Glycine max) BM085020 (Glycine max) AY059416 (Zea mays) AF273253(Beta vulgaris) L39940 (Oryza sativa) AJ316590 (Nostoc punctiforme) AJ316589 (Nostoc punctiforme) AJ316596 (Anabaena sp) AJ316595(Anabaena sp) AJ316584 (Anabaena sp) BM005654 (Crocus sativus) BI973032 (Glycine max) BI971794 (Glycine max) AF367450 (Prunus persica) BI945506 (Glycine max) BI944973 (Glycine max) AF420224 (Carica papaya) BI788449 (Glycine max) BI788359 (Glycine max) BI787127 (Glycine max) BI787033(Glycine max) BI787000 (Glycine max) BI786823(Glycine max) BI784933 (Glycine max) BI784627 (Glycine max) BI700214 (Glycine max) BI699934 (Glycine max) BI699923 (Glycine max) BI699585(Glycine max) BI543240 (Sugar beet) BI498340 (Glycine max) BI471463 (Glycine max) BI427241(Glycine max) BI427174 (Glycine max) BI427022 (Glycine max) BI426915 (Glycine max)AF393809 (Apium graveolens) BI321173 (Glycine max) BI320832(Glycine max) BI316894 (Glycine max) BI316826 (Glycine max) BI316405 (Glycine max) BI315949 (Glycine max) BI203222 (Lycopersicon esculentum) BI176503 (Solanum tuberosum) BG273882 (Grape berries) AY034958 (Arabidopsis thaliana) AF378187(Oryza sativa) BG790580 (Glycine max) BG790079 (Glycine max) BG726150 (Glycine max) BG654021 (Glycine max) BG653916 (Glycine max) BG653624 (Glycine max) BG652711(Glycine max) BG652507 (Glycine max) BG649914 (Glycine max) BG649831 (Glycine max) AJ309093 (Pinus pinaster) BG507656 (Glycine max) BG405632 (Glycine max) BG405474 (Glycine max) BG405204 (Glycine max) BG405017 (Glycine max) BG363730 (Glycine max) BG362638 (Glycine max) BG359764 (Glycine max) BG359452 (Glycine max) BG359307 (Glycine max) AJ311496 (Pisum sativum) BG239317 (Glycine max)

- BG237287 (Glycine max) BG157592 (Glycine max) BG155900 (Glycine max)
 BG047402 (Glycine max) BG046717 (Glycine max) BG046686 (Glycine max)
 BG046043 (Glycine max) BG043243 (Glycine max) BG042159 (Glycine max)
 BG041814 (Glycine max) AB045710 (Pyrus pyrifolia) BF597682 (Glycine max)
 5 BF597330 (Glycine max) BF597258 (Glycine max) BF595837 (Glycine max)
 BF595646 (Glycine max) BF425609 (Glycine max) BF423725 (Glycine max)
 BF324657 (Glycine max) BF154308 (Solanum tuberosum) BF154187
 (Solanum tuberosum) BF154037 (Solanum tuberosum) BF153500 (Solanum
 tuberosum) BF153341 (Solanum tuberosum) BF153335 (Solanum tuberosum)
 10 BF153307 (Solanum tuberosum) BF097021 (Lycopersicon esculentum)
 BF070782 (Glycine max) BF070666 (Glycine max) BF070277 (Glycine max)
 BF070188 (Glycine max) BF068815 (Glycine max) BF071174 (Glycine max)
 BF067110 (Glycine max) BF009756 (Glycine max) BE806119 (Glycine max)
 BE805996 (Glycine max) BE805178 (Glycine max) BE800976 (Glycine max)
 15 BE800941 (Glycine max) BE800227 (Glycine max) BE611805 (Glycine max)
 BE611493 (Glycine max) BE610447 (Glycine max) BE610325 (Glycine max)
 BE609887 (Glycine max) BE609881 (Glycine max) BE607382 (Oryza sativa)
 BE607326 (Oryza sativa) BE607323 (Oryza sativa) BE556484 (Glycine max)
 BE556306 (Glycine max) BE555319 (Glycine max) AJ001117 (Triticum
 20 aestivum) AJ292758 (Anabaena variabilis) BE474178 (Glycine max) BE440931
 (Glycine max) BE440323 (Glycine max) BE347277 (Glycine max) AF263384
 (Saccharum officinarum) BE330345 (Glycine max) BE211050 (Glycine max)
 BE209707 (Glycine max) AW561929 (Gossypium hirsutum) BE040389 (Oryza
 sativa) BE040121 (Oryza sativa) BE039563 (Arabidopsis thaliana) BE034387
 25 (Mesembryanthemum crystallinum) BE033387 (Mesembryanthemum
 crystallinum) BE023957 (Glycine max) BE023630 (Glycine max) BE022760
 (Glycine max) BE021167 (Glycine max) BE020760 (Glycine max) BE020591
 (Glycine max) BE020550 (Glycine max) AW990923 (Euphorbia esula)
 AW832022 (Glycine max) AW760552 (Glycine max) AW756812 (Glycine max)
 30 AW756072 (Glycine max) AW756065 (Glycine max) AW734901 (Glycine max)
 AW707163 (Glycine max) AW706660 (Glycine max) AW706520 (Glycine max)

- AW706203 (Glycine max) AW705595 (Glycine max) AW704619 (Glycine max)
 AB025778 (Citrus unshiu) AB021745 (Citrus unshiu) AW666333 (Glycine max)
 AW666277 (Glycine max) AW666250 (Glycine max) AW620859 (Glycine max)
 AW598476 (Glycine max) AW598473 (Glycine max) AW597690 (Glycine max)
 5 AW597373 (Glycine max) AW597332 (Glycine max) AW596993 (Glycine max)
 AW570615 (Glycine max) AW570577 (Glycine max) AW570566 (Glycine max)
 AW570513 (Glycine max) AW569822 (Glycine max) AW568526 (Glycine max)
 AW568333 (Glycine max) AW509231 (Glycine max) AW472408 (Glycine max)
 AW459606 (Glycine max) AW458318 (Glycine max) X82504 (C.rubrum)
 10 AW432731 (Glycine max) AW432392 (Glycine max) AW432192 (Glycine max)
 AW397142 (Glycine max) AW397071 (Glycine max) AW309503 (Glycine max)
 AW307502 (Glycine max) AW307391 (Glycine max) AW307001 (Glycine max)
 AW306834 (Glycine max) AB018561 (Citrullus lanatus) AB029401 (Citrus
 unshiu) AB022092 (Citrus unshiu) AB022091 (Citrus unshiu) AW279073
 15 (Glycine max) AW279053 (Glycine max) AW278487 (Glycine max) AJ388994
 (Medicago truncatula) AJ388888 (Medicago truncatula) AW234887 (Glycine
 max) AJ238219 (Triticum aestivum) AJ238218 (Triticum aestivum) AJ238217
 (Triticum speltoides) AW201670 (Glycine max) AW185801 (Glycine max)
 AW185627 (Glycine max) AJ249624 (Triticum aestivum) AJ249623 (Triticum
 20 aestivum) AW164630 (Glycine max) AW164393 (Glycine max) AW133248
 (Glycine max) AW101578 (Glycine max) AW100191 (Glycine max) AW100069
 (Glycine max) AW099557 (Glycine max) X96938 (T.gesneriana) X96939
 (T.gesneriana) AW035186 (Lycopersicon esculentum) AW033439
 (Lycopersicon esculentum) AW032339 (Lycopersicon esculentum) AJ132002
 25 (Craterostigma plantagineum) AJ132001 (Craterostigma plantagineum)
 AJ132000 (Craterostigma plantagineum) AJ131999 (Craterostigma
 plantagineum) AI973811 (Glycine max) AI973710 (Glycine max) AI973540
 (Glycine max) AI967739 (Lotus japonicus) AI965972 (Glycine max) AI960742
 (Glycine max) AI960703 (Glycine max) AI930917 (Glycine max) AI900130
 30 (Glycine max) AI900087 (Glycine max) AI855470 (Glycine max) AA080634
 (Saccharum sp.) AA080610 (Saccharum sp.) AA269294 (Saccharum sp.)

AA080580 (*Saccharum* sp.) AI736370 (*Glycine* max) AI731292 (*Gossypium*
hirsutum) AI731115 (*Gossypium* *hirsutum*) AI729201 (*Gossypium* *hirsutum*)
 AI728436 (*Gossypium* *hirsutum*) AI727966 (*Gossypium* *hirsutum*) AI726092
 (*Gossypium* *hirsutum*) U73588 (*Gossypium* *hirsutum*) U73587 (*Gossypium*
 5 *hirsutum*) AJ012080 (*Pisum* *sativum*) AJ131964 (*Medicago* *truncatula*)
 AJ131943 (*Medicago* *truncatula*) AJ133726 (*Lotus* *japonicus*) Y16091 (*Daucus*
carota) Y16090 (*Daucus* *carota*) AJ011319 (*Lycopersicum* *esculentum*)
 AI496671 (*Glycine* max) AI496540 (*Glycine* max) AI496532 (*Glycine* max)
 AI495774 (*Glycine* max) AI495135 (*Glycine* max) AI495023 (*Glycine* max)
 10 AI494833 (*Glycine* max) AJ011534 (*Lycopersicon* *esculentum*) Y15802
 (*Hordeum* *vulgare*) AI461126(*Glycine* max) AI460757 (*Glycine* max) AI460629
 (*Glycine* max) AI444096 (*Glycine* max) AI444083 (*Glycine* max) AI444054
 (*Glycine* max) AI443620 (*Glycine* max) AI443476 (*Glycine* max) AI443231
 (*Glycine* max) AI442789 (*Glycine* max) AI442411 (*Glycine* max) AI441989
 15 (*Glycine* max) AI441004 (*Glycine* max) AI437923 (*Glycine* max) AI437907
 (*Glycine* max) AI437840 (*Glycine* max) AJ010639 (*Anabaena* sp.) AJ011535
 (*Lycopersicon* *esculentum*) D10266 (*Vigna* *radiata*) L03366 (*Oryza* *sativa*)
 AF030231 (*Glycine* max) M97551 (*Vicia* *faba*) AJ000153 (*Triticum* *aestivum*)
 AF079523 (*Musa* *acuminata*) AF079851 (*Pisum* *sativum*) AJ001071 (*Pisum*
 20 *sativum*) AF049487 (*Medicago* *sativa*) AF054446 (*Mesembryanthemum*
crystallinum) AA753339 (*Oryza* *sativa*) AA752298 (*Oryza* *sativa*) AA752293
 (*Oryza* *sativa*) AA753445 (*Oryza* *sativa*) AA753437 (*Oryza* *sativa*) AA753297
 (*Oryza* *sativa*) AA752123 (*Oryza* *sativa*) AA751990 (*Oryza* *sativa*) AA750692
 (*Oryza* *sativa*) AA750079 (*Oryza* *sativa*) AA749692 (*Oryza* *sativa*) AA749554
 25 (*Oryza* *sativa*) AA720478 (*Mesembryanthemum* *crystallinum*) AA661050
 (*Medicago* *truncatula*) AA661041 (*Medicago* *truncatula*) AA660686 (*Medicago*
truncatula) D88412 (*Cotton*) D10418 (*Rice*) D21308 (*Rice*) D29733 (*Rice*)
 X81974 (*B.vulgaris*) X92378 (*A.glutinosa*) Z56278 (*V.faba*) Z48640 (*V.faba*)
 X98598 (*P.sativum*) T25261 (*Zea* *mays*) T23326 (*Zea* *mays*) T14713 (*Zea*
 30 *mays*) T14662 (*Zea* *mays*) T14661 (*Zea* *mays*) X75332 (*D.carota*) X02382
 (*Zea* *mays*) X02400 (*Zea* *mays*) X70990 (*A.thaliana*) X60987 (*A.thaliana*)

X69773 (V.faba) X73477 (S.tuberosum) Z11532 (S.officinatum) Z15028 (O.sativa) X64770 (O.sativa) X59046 (O.sativa) X66728 (H.vulgare) X65871 (H.vulgare) X69931 (H.vulgare) A27685 (O.sativa) W21612 (Zea mays) U24088 (Solanum tuberosum) U24087 (Solanum tuberosum) X73221 (H.vulgare)
5 L32898 (Zea mays) F13913 (Arabidopsis thaliana) F13912 (Arabidopsis thaliana) U21129 (Solanum tuberosum) M26672 (Triticum aestivum) M26671 (Triticum aestivum) L19762 (Lycopersicon esculentum) M18745 (Potato) L33244 (Zea mays) L22296 (Zea mays) Z17959 (Arabidopsis thaliana). These sequences are incorporated by reference.

10

In a second aspect, the present invention provides a method of altering the production of fibre, fruit or seeds in a plant, the method comprising causing under- or overexpression of an iso-form of SuSy involved in fruit/seed production in the plant.

15

Preferably, for underexpression, the method involves providing a genetic construct which targets the 3' end of the SuSy gene when transformed in a plant. By "3'-end" is meant a sequence encoding the C-terminal portion of a SuSy polypeptide with or without the 3' non-coding sequence, or a sequence
20 that is complementary thereto. The construct can be a co-suppression antisense or combined sense/antisense (inverted repeat) SuSy construct.

In a third aspect, the present invention provides a plant having an altered ability to produce fibre, fruit or seeds produced by the method according to the
25 second aspect of the present invention.

In a fourth aspect, the present invention provides a genetic construct targeting the 3' end of the SuSy gene which, when expressed in a plant, reduces the expression of an iso-form of SuSy involved in fruit/seed production.

30

Preferred embodiments of the chimeric genes, particularly of the coding regions used for these methods are as described elsewhere in this specification.

Preferably, the construct is selected from the constructs shown in Figure 5, a
5 combination of parts of both constructs to generate an inverted repeat gene suppression construct, with or without a tissue specific promoter region.

One preferred method of developing constructs is by using the "pPLEX" technology involving subclover stunt virus promoters and terminators described
10 in AU 689311.

In a fifth aspect, the present invention provides use for of the construct according to the fourth aspect of the present invention to produce a plant having reduced ability to produce fibre, fruit or seeds.

15

The present invention is suitable for the following applications:-

A) Use of SuSy to enhance or otherwise improve fibre yield

-SuSy may be used to enhance or otherwise improve fibre synthesis by increasing SuSy expression or activity in such plant cells e.g. a fibre specific or
20 fibre enhanced promoter (such as but not limited to a cotton expansion promoter, or the promoter of the SuSy gene having the nucleotide sequence of SEQ ID NO: 1) or a constitutive promoter (such as CaMV 35S) can be used to express the SuSy gene in cotton cells.

-Cotton fibre yield can be improved by reducing or suppressing cotton
25 fuzz fibre. Fuzz fibre specific or enhanced promoter or constitutive promoter to reduce sucrose synthase selectively in cotton fuzz fibre cells. Reduced expression can be achieved by gene silencing technologies (antisense, co-suppression, dominant negative etc). Dominant negative mutant alleles of sucrose synthase genes may be obtained e.g. through mutation (insertion,
30 substitution, or deletion of the phosphorylation site(s) of the sucrose synthase protein coding region)

- Cotton fibre yield can be improved by modifying fuzz fibres into lint fibres e.g. by expression of SuSy coding region or a functional equivalent thereof by fibre specific or fibre enhanced promoter, primary or secondary cell wall promoter or a constitutive promoter.

5

B) Use of SuSy gene to improve fibre quality

-Fibre length is an example of fibre quality. Expression of SuSy by a fibre specific or fibre enhanced promoter, primary cell wall promoter or a constitutive promoter.

10

C) Use of SuSy to improve fibre properties (e.g. fibre strength, length and number)

-Fibre strength is an example of a fibre property. Fibre strength is significantly effected by the cellulose content in the secondary cell wall of fibre cells. This may be achieved by expression of SuSy by a fibre specific or fibre enhanced promoter, secondary cell wall promoter or a constitutive promoter.

15

D) Overexpression of SuSy specifically in seeds to increase sucrose utilisation in seeds for increased seed size and storage product content.

20

E) Overexpression in the maternal tissue of a fruit (such as cotton fibre, horticultural fruit) for increased carbohydrate or fibre content.

F) Suppression of SuSy in seed with over expression of SuSy in fruit/fibre for increased fibre/fruit yield.

25

Preferred plant-operable promoters include the fibre specific and/or secondary cell wall specific promoters which can be isolated according to the teaching of WO 98/18949, WO98/00549 or US5932713.

30

The correlation of sucrose synthase activity in the ovule epidermis of cotton with fibre length also allows the use of sucrose synthase proteins (or antibodies or aptamers recognizing the same) or sucrose synthase coding regions as a diagnostic tool in cotton breeding. These may be used to identify cotton lines or varieties, including wild sources, which have enhanced sucrose synthase activity and increased potential to form longer fibres, particularly when crossed in a breeding program with other cotton lines and/or varieties having good fibre characteristics. Nucleic acids derived from sucrose synthase coding regions may be used to determine the amount of sucrose synthase RNA in the ovule epidermis. Also, polymorphism, including single nucleotide polymorphism between sucrose synthase genes may be used to identify lines with superior sucrose synthase alleles.

The methods and means of the invention are particularly suited for use in cotton plants, (both *Gossypium hirsutum* and *Gossypium barbadense*) particularly for Coker 312, Coker310, Coker 5Acala SJ-5, GSC25110, FiberMax 819, Siokra 1-3, T25, GSA75, Acala SJ2, Acala SJ4, Acala SJ5, Acala SJ-C1, Acala B1644, Acala B1654-26, Acala B1654-43, Acala B3991, Acala GC356, Acala GC510, Acala GAM1, Acala C1, Acala Royale, Acala Maxxa, Acala Prema, Acala B638, Acala B1810, Acala B2724, Acala B4894, Acala B5002, non Acala "picker" Siokra, "stripper" variety FC2017, Coker 315, STONEVILLE 506, STONEVILLE 825, DP50, DP61, DP90, DP77, DES119, McN235, HBX87, HBX191, HBX107, FC 3027, CHEMBRED A1, CHEMBRED A2, CHEMBRED A3, CHEMBRED A4, CHEMBRED B1, CHEMBRED B2, CHEMBRED B3, CHEMBRED C1, CHEMBRED C2, CHEMBRED C3, CHEMBRED C4, PAYMASTER 145, HS26, HS46, SICALA, PIMA S6 and ORO BLANCO PIMA.

Any description of prior art documents herein is not an admission that the documents form part of the common general knowledge of the relevant art in Australia.

In order that the present invention may be more clearly understood preferred forms will be described with reference to the following examples and accompanying drawings.

5 Modes for Carrying Out the Invention

General molecular biology

Unless otherwise indicated, the recombinant DNA techniques utilised in the present invention are standard procedures, well known to those skilled in the art. Such techniques are described and explained throughout the literature in
10 sources such as, J. Perbal, A Practical Guide to Molecular Cloning, John Wiley and Sons (1984), J. Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Cold Spring Harbour Laboratory Press (1989), T.A. Brown (editor), Essential Molecular Biology: A Practical Approach, Volumes 1 and 2, IRL Press (1991),
15 D.M. Glover and B.D. Hames (editors), DNA Cloning: A Practical Approach, Volumes 1-4, IRL Press (1995 and 1996), and F.M. Ausubel *et al.* (Editors), Current Protocols in Molecular Biology, Greene Pub. Associates and Wiley-Interscience (1988, including all updates until present) and are incorporated herein by reference.

20 Analysis for presence of the transgene by Southern blotting was carried out as described in Sambrook *et al.* (1989) and immunolocalisation of the SuSy protein was carried out on tissue sections as described in Ruan and Chourey (1998).

25 Gene/DNA isolation

The DNA encoding a protein may be obtained from any cDNA library prepared from tissue or organisms believed to express the gene mRNA and to express it at a detectable level. The gene sequences can also be obtained from a genomic library or genomic DNA.

30

- Libraries are screened with probes or analytical tools designed to identify the gene of interest or the protein encoded by it. For cDNA expression libraries, suitable probes include monoclonal or polyclonal antibodies that recognise and specifically bind the protein; oligonucleotides of about 20-80 bases in length
- 5 that encode known or suspected portions of cDNA from the same or different species; and/or complementary or homologous cDNAs or fragments thereof that encode the same or a hybridising gene. Appropriate probes for screening genomic DNA libraries include, but are not limited to, oligonucleotides; cDNAs or fragments thereof that encode the same or hybridising DNA including
- 10 expressed sequence tags and the like; and/or homologous genomic DNAs or fragments thereof. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures as described in chapters 10-12 of Sambrook *et al.*
- 15 An alternative means to isolate a gene encoding is to use polymerase chain reaction (PCR) methodology as described in section 14 of Sambrook *et al.* This method requires the use of oligonucleotide probes that will hybridise to the gene.
- 20 The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimised. The actual nucleotide sequence(s) is usually based on conserved or highly homologous nucleotide sequences or regions of the gene. The oligonucleotides may be degenerate at one or more positions. The use of
- 25 degenerate oligonucleotides may be of particular importance where a library is screened from a species in which preferential codon usage in that species is known. The oligonucleotide must be labelled such that it can be detected upon hybridisation to DNA in the library being screened. The preferred method of labelling is to use ^{32}P -labelled dATP with polynucleotide kinase, as is well
- 30 known in the art, to radiolabel the oligonucleotide. However, other methods

may be used to label the oligonucleotide, including, but not limited to, biotinylation or enzyme labelling.

5 Nucleic acid having all the protein coding sequence is obtained by screening selected cDNA or genomic libraries, and if necessary, using conventional primer extension procedures as described in section 7.79 of Sambrook *et al.*, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

10 Another alternative method for obtaining the gene of interest is to chemically synthesise it using one of the methods described in Fingels *et al.* (Agnew Chem. Int. Ed. Engl. 28: 716-734, 1989). These methods include triester, phosphite, phosphoramidite and H-Phosphonate methods, PCR and other autoprimer methods, and oligonucleotide syntheses on solid supports. These
15 methods may be used if the entire nucleic acid sequence of the gene is known, or the sequence of the nucleic acid complementary to the coding strand is available, or alternatively, if the target amino acid sequence is known, one may infer potential nucleic acid sequences using known and preferred coding residues for each amino acid residue.

20

Cotton Transformation

Cotton was transformed using an *Agrobacterium* mediated transformation technique as described in F. Murray *et al.* 1999 (Molec. Breeding 5: 219-232). Transformed plants were grown in a glasshouse under natural illumination at
25 day/night temperatures of approximately 30°C / 22°C.

Example 1: Suppression of SuSy expression in cotton.

To provide evidence regarding the role of SuSy in fibre development, the present inventors transformed cotton with co-suppression and antisense SuSy
30 constructs, targeting the 3' end of the seed SuSy cDNA driven by a constitutive subclover stunt seven virus promoter (see Figure 5 for a schematic

representation). The presence of the transgene in 11 transgenic lines so far was confirmed by Southern analysis (see Figure 6). Among them two antisense lines (294-147, 292-43) and one co-suppression line (295-82) showed dramatic reductions in fibre and seed development with the remaining lines showing various degrees of inhibition of fibre growth.

Immunolocalization analysis on 0-d ovule sections (Figure 2) revealed that, compared to wild type fibre initials, SuSy protein was reduced to less than 20% wild type levels in fibres of line 294-147 and to undetectable levels in 295-82. Furthermore, the number and size of initiating fibres was reduced by at least 50% in these two lines as compared that in the wild type ovules. Indeed, by 6 days after anthesis (DAA), these transgenic seeds were virtually fibre-less (see Figure 1). Sucrose synthase activity was determined biochemically. Whereas WT plants show a SuSy activity of 22.5 nmol/min/seed, line 147 showed a SuSy activity of 5.8 nmol/min/seed and line 82 showed no detectable activity. This led to a dramatic reduction of normal fibre initial cells in the ovule epidermis. It is clear from these results that the degree of fibre initiation depends on the level of SuSy expression in the ovule epidermis.

The impact of SuSy expression on fibre initiation is further demonstrated by scanning electron microscopy analysis of the ovule epidermis of line 147 (see figure 7) revealed that many fibre cells are collapsed and shrunk (see figure 7B). Some fibre cells appeared, but these were much smaller in size, and were reduced in number when compared to wild type ovule (figure 7A). The shrunk and collapsed phenotype was clearly visible under high magnification (figure 7C).

These results demonstrate that some iso-forms of SuSy play a critical role in fibre initiation and elongation. Suppression of an iso-form of SuSy can thus be used to prevent or reduce fibre initiation and elongation while overexpression can thus be used to enhance fibre initiation and elongation.

While all the bolls in line 295-82 and 292-43 dropped off prematurely by 10 DAA, some bolls of line 294-147 were retained to maturity. In those mature bolls, most seeds were stunted, shrunken and fibre-less. About 15% of the seeds, however, showed 30% of the wild type fibre length (loose fibre as well) and wild type embryo and seed size, most likely due to segregation of the transgene.

This demonstrates that suppression of iso-forms of SuSy in the maternal tissue (seed coat/ fibre) alone can inhibit fibre development by ~70% in length, while additional repression of SuSy in the embryo can arrest seed development entirely.

Gene suppression of iso-forms of SuSy in the embryo specifically to give a seedless phenotype (in a wide range of horticultural crops such as grape, peach, pear, apple).

Line 292-43 and 295-82 were male-sterile (no pollen) and the seed set can be recovered by pollination with wild-type pollen. This shows SuSy plays a role in male sterility of cotton.

Gene suppression of SuSy in male floral parts was found to cause sterility. Sucrose synthase activity, the fresh weight of pollen and viability of the pollen were tested on developing anthers 2 days before flowering in different lines. The viability assay was conducted by staining with 2,3,5-triphenyltetrazolium chloride (TTC). The results are summarized in Table 1.

Table 1. Suppression of SuSy in anthers leads to male sterility by pollen formation inhibition.

Line	SuSy activity (nmol min ⁻¹ mg ⁻¹)	Pollen fresh weight (% of WT pollen per anther)	Viable pollen (% of 100 grains)
WT	44.1 ± 1.1	100.0	85.2
147#1	29.5 ± 6.9	30.0	52
147#2	14.7 ± 0.9	0.0	0.0
82	18.2 ± 1.5	0.0	0.0

A reduction of SuSy activity in anthers by about 33% results in an about 70%
 5 reduction of total pollen weight, largely due to the reduction of the number of
 pollen produced. Of the residual 30%, only half of the pollen was viable. When
 SuSy was reduced by about 60% no pollen was any longer formed. Pollen
 development is thus very sensitive to changes in SuSy activity. Pollen
 formation can thus be influenced (inhibited or promoted) by suppressing or
 10 overexpressing SuSy in anthers.

Suppression of SuSy activity in the maternal seed coat tissue in the transgenic
 cotton plants, also led to reduced fuzz length in the SuSy suppressed
 transgenic plants when compared to a wild type plant (see Figure 9). Fuzz fibre
 15 length was measured from the chalazal end of the mature seeds and the
 measurement was done by gently stretching the fuzz of seeds, delimited by
 hand, and measuring the length from the seed coat epidermis to the tip. The
 brown color visible in the transgenic line, is due to the fact that fuzz is shorter in
 transgenic lines than in WT, where the brown seed coat is well covered by
 20 relatively long fuzz. SuSy activity was also determined. The results can be
 summarized as follows:

- the wild type plant had a fuzz length of 2.2 mm (and a SuSy activity measurement set at 100%)

- transgenic line 147#2 had a fuzz length of 1.4 mm (and a SuSy activity measurement of 36% of the WT).

From the general correlation detected between SuSy expression and fibre
5 content and/or quality described herein, it will be understood that
overexpression of SuSy gene will have potential beneficial effects on fibre, fruit
and seed production in plants.

Example 2. Analysis of progeny of cotton lines 82, 147 and 43 with suppressed
10 SuSy activity.

To confirm the correlation between the presence of the SuSy suppression
transgene and the reduced fibre initiation in cotton, further analysis was
conducted on the T1 generation. About 24 T1 seeds were sown from 6 T0
lines. All germinated seedlings were screened by PCR for presence or absence
15 of the transgenes (nptII; S7 promoter and SuSy suppression transgene). Six
PCR positive and one PCR negative segregating individuals from each of the 6
lines were further analysed on molecular biochemical and cellular level. Line 82
was propagated vegetatively as a reference point. The fibre-less phenotype in
the different lines of the T0 generation (indicated by arrows in figure 6A) was
20 still preserved in respectively 3 and 2 segregating individuals of the T1
generation for line 147 and 43 respectively (arrows in Figure 6B). The
remaining individuals show various degree of fibre/ seed suppression. The
vegetatively-kept line 82 continues to produced fibreless seeds. These results
demonstrated that the fibreless/ or seed suppressed phenotype described was
25 indeed due to the presence of SuSy transgene.

A positive strong correlation could be found between the degree of sucrose
activity in the seed epidermis and the fibre length in segregating T1 generation
plants of lines 147, 43 and 101 (see Figures 8, 9 and 10). When fibre length
30 was plotted against sucrose synthase activity for the different plants of the
different lines as well as the wild type line, a linear regression could be derived

($Y=2.634X-50$; $R^2=0.98$). The linear nature of the correlation up to the wild-type level, indicates that overexpression of the sucrose synthase activity above wild type level will lead to fibre length greater than fibre length in wild type plants.

- 5 A similar strong correlation exists between the reduction in SuSy activity and mature cotton fibre length and dry weight. SuSy activity in seed coat epidermis has been reduced to 28.8%, 42.5% and 53.9% of the wild type level in segregating individuals #1, #2 and #3 of line 43 T1 generation plants respectively. Lint fibre dry weight per boll was 11.1%, 51.6%, and 70.9%
10 respectively.

Cellulose content in cotton fibre is also closely correlated to the level of SuSy expression. Cellulose content was analysed by labelling cotton fibre at day 25 (peak of cellulose synthesis *in vivo*) with Calcofluor white, a fluorescent dye
15 specifically binding to cellulose. The fluorescent intensity of the labelled fibres, indicating cellulose content, was significantly reduced in the SuSy suppressed transgenic lines when compared to fluorescence in the wild type. The degree of fluorescence reduction correlated to the degree of suppression of SuSy activity.

20 **Example 3. Overexpression of sucrose synthase in cotton plants.**

The coding sequence of a potato sucrose synthase cDNA (Genbank Accession number M18745) is operably linked to a subterranean clover stunt virus promoter (S7; WO9606932) and a 3' transcription termination and polyadenylation signal functional in plants. This chimeric gene is operably
25 linked to a selectable marker gene and introduced into a T-DNA vector. Cotton plants are transformed using the above mentioned Agrobacterium mediated transformation technique. Transgenic cotton lines are identified, sucrose synthase activity, fibre length, fuzz fibre length, cellulose content, and dry weight of the lint is analyzed. A positive correlation is found between SuSy
30 activity and increased fibre length cellulose content, and dry weight of the lint.

SUMMARY OF THE EXAMPLES

The single-cell cotton fibres initiate from ovule epidermis at anthesis, elongate to 2.5 ~ 3.0 cm in about 16 days and then synthesize massive amounts of cellulose. Thus, cotton fibre is an excellent system for the study of cell
5 differentiation, elongation and cellulose synthesis in higher plants with significant industry implications for improving fibre yield and quality.

To provide definitive evidence regarding the role of SuSy in fibre and seed development, the present inventors transformed cotton with co-suppression
10 and antisense SuSy constructs, targeting the 3' end of the seed SuSy cDNA driven by subclover stunt seven virus promoter. The presence of the transgene in 9 transgenic lines so far was confirmed by Southern analysis. Among them one antisense line (294-147) and one co-suppression line (295-82) showed dramatic reductions in fibre and seed development with the remaining lines
15 showing various degrees of inhibition of fibre growth. Immunolocalization analysis on 0-d ovule sections revealed that, compared to wild type fibre initials, SuSy protein was reduced to only about 20% wild type levels in fibres of line 294-147 and to undetectable levels in 295-82. Furthermore, the number and size of initiating fibres was reduced by at least 50% in these two lines as
20 compared that in the wild type ovules. Indeed, by 2 d after anthesis (DAA), these transgenic seeds were virtually fibre-less. This is in contrast to a fibre-covered seed phenotype seen in the wild type plants at this stage. While all the bolls in line 295-82 dropped off prematurely by 10 DAA, some bolls of line 294-147 were retained to maturity. In those mature bolls, most seeds were stunted,
25 shrunken and fibre-less. About 15% of the seeds, however, showed 30% of the wild type fibre length and wild type embryo and seed size, most likely due to segregation of the transgene. These results demonstrate (a) that SuSy plays a critical role in fibre initiation and elongation, (b) that suppression of SuSy in the maternal tissue (seed coat/fibre) alone can inhibit fibre
30 development while additional repression of SuSy in the embryo can arrest seed development entirely. The influence of SuSy suppression on the reduction of

fibre length has been confirmed by analysis of progeny of the transgenic cotton lines. Further, a linear correlation was found between the level of SuSy activity and the increase in fibre length. The linear nature of this correlation indicates that in wild type cotton, the level of sucrose synthase is limiting.

5

Overexpression of sucrose synthase at least in maternal ovule tissue increases fibre length in fibre producing plants.

10 It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

References:

15 Amor , Y., Haigler, C.H., Johnson, S., Wainscott, M. and Delmer, D.P. (1995) A membrane associated form of sucrose synthase and its potential role in synthesis of cellulose and callose in plants. *Proc Nat Acad Science USA* 92, 9353-9357.

20 Ruan, Y.-L., Llewellyn, D.J, Furbank, R.T. (2000) Pathway and control of sucrose import into initiating fibre cells. *Australian Journal of Plant Physiology* 27, 795-800.

25 Ruan, Y.-L. and Chourey, P.S. (1998) A fibreless seed (fls) mutation in cotton is associated with lack of fibre cell initiation in ovule epidermis, alteration in sucrose synthase expression and carbon partitioning in developing seed. *Plant Physiology* 118: 399-406

30 Ruan, Y.-L., Chourey, P.S., Delmer, D.P. and Luis, P.G. (1997) The differential expression of sucrose synthase in relation to diverse patterns of carbon partitioning in developing cotton seed. *Plant Physiology* 115: 375-385

We claim:

1. A method for altering fibre development or properties of a fibre producing plant comprising the steps of
 - a) providing cells of said plants with a chimeric gene comprising the following operably linked DNA fragments:
 - (i) a plant operable promoter;
 - (ii) a coding region as hereinbefore defined which when transcribed yields an RNA said RNA being capable of reducing the expression of an endogenous sucrose synthase gene or capable of being translated into an active sucrose synthase protein; and
 - (iii) transcription termination and polyadenylation signals that function in said plant cells.
2. The method of claim 1 wherein the endogenous sucrose synthase gene is a gene that is expressed in fibre cells.
3. The method of claim 2 wherein the fibre cells are fibre initial cells.
4. The method according to any one of claims 1 to 3, wherein said RNA is capable of being translated into an active sucrose synthase protein.
5. The method according to claim 4, wherein said coding region comprises a nucleotide sequence selected from the group consisting of:
 - a) a nucleotide sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2 or a fragment of said nucleotide sequence encoding an active sucrose synthase;

- b) the nucleotide sequence of SEQ ID NO: 1 or a fragment of said nucleotide sequence encoding an active sucrose synthase;
 - c) a nucleotide sequence having at least 70% sequence identity to a) or b);
 - d) a nucleotide sequence that hybridizes under stringent conditions with a) or b) or a complementary sequence thereto or a part thereof encoding an active sucrose synthase.
6. The method according to any one of claims 1 to 3, wherein said coding region is transcribed to produce RNA that is capable of reducing the expression of an endogenous sucrose synthase gene.
7. The method according to claim 6, wherein said coding region comprises a nucleotide sequence selected from the group consisting of:
- a) a nucleotide sequence comprising at least about 19 contiguous nucleotides of a sequence that has at least 70% sequence identity to a nucleotide sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2;
 - b) a nucleotide sequence comprising at least about 25 contiguous nucleotides of a sequence having at least 70% sequence identity to the nucleotide sequence of SEQ ID NO: 1;
 - c) the complement of a) or b);
8. The method of claim 7 wherein the nucleotide sequence at a) comprises at least about 25 contiguous nucleotides of a sequence that has at least 70% sequence identity to a nucleotide sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2.
9. The method according to any one of claims 6 to 8, wherein said coding region comprises the nucleotide sequence of SEQ ID NO: 1 from the

nucleotide at about position 2208 to the nucleotide at about position 2598 or the complement thereof.

10. The method according to any one of claims 6 to 9, wherein said coding region comprises both sense and antisense nucleotide sequences capable of forming a double stranded RNA molecule.
11. The method according to any one of claims 1 to 10, wherein said promoter is a subterranean clover stunt virus promoter.
12. The method according to any one of claims 1 to 11, wherein said fibre producing plant is a cotton plant.
13. The method according to claim 12, wherein said cotton plant is a FibermaxTM variety.
14. A method for enhancing fibre yield in a fibre producing plant comprising the steps of:
 - a) providing cells of said plant with a chimeric gene comprising the following operably linked DNA fragments:
 - i) a plant operable promoter;
 - ii) a DNA region capable of being translated into an active sucrose synthase protein; and
 - iii) transcription termination and polyadenylation signals that function in said plant cells.

15. A method for enhancing fibre quality in a fibre producing plant, comprising the steps of
 - b) providing cells of said plant with a chimeric gene comprising the following operably linked DNA fragments
 - i) a plant operable promoter;
 - ii) a DNA region capable of being translated into an active sucrose synthase protein
 - iii) transcription termination and polyadenylation signals that function in said plant cells.
16. A method for increasing seed size in a fibre producing plant, comprising the steps of:
 - a) providing cells of said plant with a chimeric gene comprising the following operably linked DNA fragments
 - b) a seed-specific promoter;
 - c) a DNA region capable of being translated into an active sucrose synthase protein; and
 - d) transcription termination and polyadenylation signals that function in said plant cells.
17. A fibre producing plant comprising in its genome a chimeric DNA comprising the following operably linked DNA fragments:
 - i) a plant operable promoter;
 - ii) a coding region which when transcribed yields an RNA said RNA being capable of reducing the expression of an endogenous

sucrose synthase gene or capable of being translated into an active sucrose synthase protein; and


- iii) transcription termination and polyadenylation signals that function in said plant cells.
18. The fibre producing plant of claim 17 wherein the coding region is transcribed to yield RNA that is capable of reducing the expression of an endogenous sucrose synthase gene in fibre cells.
 19. The fibre producing plant of claim 18 wherein the fibre cells are fibre initial cells.
 20. The fibre producing plant according to claim 17, wherein said RNA is translated into an active sucrose synthase protein.
 21. The fibre producing plant of claim 20 wherein the active sucrose synthase protein is expressed in fibre cells and wherein said fibre cells have an increased sucrose synthase activity compared to fibre cells of plant cells which do not comprise said chimeric DNA.
 22. The fibre producing plant of claim 21 wherein the fibre cells are fibre initial cells.

23. The fibre producing plant according to any one of claims 17 to 22, wherein that coding region comprises a nucleotide sequence selected from the group consisting of:
- a) a nucleotide sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2 or a fragment of said nucleotide sequence encoding an active sucrose synthase;
 - b) the nucleotide sequence of SEQ ID NO: 1 or a fragment of said nucleotide sequence encoding an active sucrose synthase;
 - c) a nucleotide sequence having at least 70% sequence identity to a) or b); and
 - d) a nucleotide sequence that hybridizes under stringent conditions with a) or b) or a complementary sequence thereto or a part thereof encoding an active sucrose synthase.
24. The fibre producing plant according to claim 23, wherein said coding region comprises a nucleotide sequence selected from the group consisting of
- a) a nucleotide sequence comprising at least about 19 contiguous nucleotides of a sequence that has at least 70% sequence identity to a nucleotide sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2;
 - b) a nucleotide sequence comprising at least about 25 contiguous nucleotides of a sequence having at least 70% sequence identity to the nucleotide sequence of SEQ ID NO: 1;
 - c) the complement of a) or b);
25. The fibre producing plant of claim 24 wherein the nucleotide sequence at a) comprises at least about 25 contiguous nucleotides of a sequence that

has at least 70% sequence identity to a nucleotide sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2.

26. The fibre producing plant according to any one of claims 17 to 25, wherein said coding region comprises the nucleotide sequence of SEQ ID No: 1 from the nucleotide at about position 2208 to the nucleotide at about position 2598 or the complement thereof.
27. The fibre producing plant according to any one of claims 17 to 26, wherein said plant is a cotton plant.
28. Seeds of the plant according to any one of claims 17 to 27.
29. Fibre is olated from the plant according to any one of claims 17 to 27 or the seed of claim 28, wherein said fibre has altered development or an altered property relative to the fibre from a plant that does not comprise the chimeric DNA.
30. A plant obtained by performing the method according to any one of claims 1 to 15.

1/11



2/11




FIGURE 2

3/11




FIGURE 3

4/11



FIGURE 4

5/11




FIGURE 5

6/11






FIGURE 6

7/11

14



14



WT




FIGURE 7

8/11




FIGURE 8

9/11




FIGURE 9

10/11




FIGURE 10

11/11




FIGURE 11

SEQUENCE LISTING

5 <110> Commonwealth Scientific Industrial and Research Organisation
 <120> Modification of sucrose synthase gene expression in plant tissue and
 uses therefor

10 <130> 500159/MRO
 <150> PR1975
 <151> 2000-12-08

15 <150> US60/251852
 <151> 2000-12-08
 <160> 2

20 <170> PatentIn version 3.1
 <210> 1
 <211> 2625
 <212> DNA
 25 <213> Gossypium hirsutum
 <220>
 <221> misc feature
 <222> (1240)..(1240)
 30 <223> N is any nucleotide
 <220>
 <221> exon
 35 <222> (1)..(2418)
 <223>

<400> 1
 40 atg gct gag cgt gct ctc act cgc gtc cac agt ctc cgt gag cgt ttg 48
 Met ala Glu Arg Ala Leu Thr Arg Val His Ser Leu Arg Glu Arg Leu
 1 5 10 15
 gat gag acc ctt ctt gct cac agg aac gag att ttg gcc ttg ctc tca 96
 45 Asp Glu Thr Leu Leu Ala His Arg Asn Glu Ile Leu Ala Leu Leu Ser
 20 25 30
 agg atc gag ggc aaa gga aaa gga att ctg caa cac cat caa att att 144
 50 Arg Ile Glu Gly Lys Gly Lys Gly Ile Leu Gln His His Gln Ile Ile
 35 40 45
 cta gag ttt gaa gct atc cct gaa gag aac aga aag aag ctc gct aat 192
 Leu Glu Phe Glu Ala Ile Pro Glu Glu Asn Arg Lys Lys Leu Ala Asn
 50 55 60
 55 ggt gca ttt ttt gaa gta ttg aag gct agt cag gaa gcg atc gtg ttg 240
 Gly Ala Phe Phe Glu Val Leu Lys Ala Ser Gln Glu Ala Ile Val Leu
 65 70 75 80
 60 cct cca tgg gtt gca ctt gct gtt cgt cca agg cct ggt gtt tgg gag 288
 Pro Pro Trp Val Ala Leu Ala Val Arg Pro Arg Pro Gly Val Trp Glu
 85 90 95
 tac att aga gtg aat gtt cac gcc ctt gtt gtt gag gaa ctc act gtt 336

	Tyr	Ile	Arg	Val	Asn	Val	His	Ala	Leu	Val	Val	Glu	Glu	Leu	Thr	Val	
				100					105					110			
5	gct	gag	tat	ctc	cac	ttc	aag	gaa	gag	ctt	gtt	gat	gga	agt	tca	aat	384
	Ala	Glu	Tyr	Leu	His	Phe	Lys	Glu	Glu	Leu	Val	Asp	Gly	Ser	Ser	Asn	
			115					120					125				
10	gga	aac	ttt	gtt	ttg	gaa	ttg	gat	ttt	gag	ccc	ttc	aac	tca	tca	ttc	432
	Gly	Asn	Phe	Val	Leu	Glu	Leu	Asp	Phe	Glu	Pro	Phe	Asn	Ser	Ser	Phe	
			130				135					140					
15	ccc	cgc	cca	act	ctt	tca	aaa	tcc	att	ggt	aat	ggt	gtg	gag	ttc	cta	480
	Pro	Arg	Pro	Thr	Leu	Ser	Lys	Ser	Ile	Gly	Asn	Gly	Val	Glu	Phe	Leu	
						150					155					160	
20	aat	cgt	cac	ctt	tcg	gca	aaa	ttg	ttc	cat	gac	aag	gag	agc	atg	cac	528
	Asn	Arg	His	Leu	Ser	Ala	Lys	Leu	Phe	His	Asp	Lys	Glu	Ser	Met	His	
					165					170					175		
25	cct	ttg	ctc	gaa	ttc	ctc	aga	gtc	cat	tgt	cac	aag	ggc	aag	aac	atg	576
	Pro	Leu	Leu	Glu	Phe	Leu	Arg	Val	His	Cys	His	Lys	Gly	Lys	Asn	Met	
				180				185						190			
30	atg	ttg	aat	gac	aga	att	cag	aac	ttg	aat	gct	ctt	caa	cat	gtt	ttg	624
	Met	Leu	Asn	Asp	Arg	Ile	Gln	Asn	Leu	Asn	Ala	Leu	Gln	His	Val	Leu	
			195				200						205				
35	agg	aaa	gca	gag	gag	tat	ctt	ggt	acc	cta	cct	cct	gag	aca	cca	tgt	672
	Arg	Lys	Ala	Glu	Glu	Tyr	Leu	Gly	Thr	Leu	Pro	Pro	Glu	Thr	Pro	Cys	
			210				215					220					
40	gcc	gaa	ttc	gaa	cac	cgg	ttc	cag	gaa	atc	ggt	ttg	gaa	aga	ggt	tggt	720
	Ala	Glu	Phe	Glu	His	Arg	Phe	Gln	Glu	Ile	Gly	Leu	Glu	Arg	Gly	Trp	
			225			230				235						240	
45	ggt	gac	acc	gca	gaa	cgc	gtg	ctc	gag	atg	atc	caa	ctc	ctt	ttg	gat	768
	Gly	Asp	Thr	Ala	Glu	Arg	Val	Leu	Glu	Met	Ile	Gln	Leu	Leu	Leu	Asp	
				245				250							255		
50	ctt	ctt	gag	gca	act	gat	cct	tgc	acc	ctt	gag	aag	ttc	ctt	ggg	aga	816
	Leu	Leu	Glu	Ala	Thr	Asp	Pro	Cys	Thr	Leu	Glu	Lys	Phe	Leu	Gly	Arg	
				260				265						270			
55	atc	ccc	atg	gtg	ttc	aat	gtt	gtg	att	ctc	act	ccc	cac	gga	tac	ttc	864
	Ile	Pro	Met	Val	Phe	Asn	Val	Val	Ile	Leu	Thr	Pro	His	Gly	Tyr	Phe	
			275				280						285				
60	gct	caa	gac	aat	gtt	ttg	ggg	tat	ccc	gac	acc	ggt	ggg	cag	gtt	gtt	912
	Ala	Gln	Asp	Asn	Val	Leu	Gly	Tyr	Pro	Asp	Thr	Gly	Gly	Gln	Val	Val	
			290				295					300					
65	tac	atc	ttg	gat	caa	gtc	cga	gct	ttg	gag	aat	gag	atg	ctc	ctc	cgt	960
	Tyr	Ile	Leu	Asp	Gln	Val	Arg	Ala	Leu	Glu	Asn	Glu	Met	Leu	Leu	Arg	
					310						315					320	
70	ata	aag	caa	caa	gga	ctc	aac	atc	acc	cct	cga	atc	ctc	att	att	act	1008
	Ile	Lys	Gln	Gln	Gly	Leu	Asn	Ile	Thr	Pro	Arg	Ile	Leu	Ile	Ile	Thr	
					325					330					335		
75	aga	ctt	ctt	cct	gat	gct	gtc	gga	aca	aca	tgc	ggt	caa	cga	ctt	gag	1056
	Arg	Leu	Leu	Pro	Asp	Ala	Val	Gly	Thr	Thr	Cys	Gly	Gln	Arg	Leu	Glu	
				340				345					350				
80	aaa	gta	tac	gga	aca	gag	cac	tcg	gat	att	ctt	cga	gta	ccc	ttc	aga	1104

	Lys	Val	Tyr	Gly	Thr	Glu	His	Ser	Asp	Ile	Leu	Arg	Val	Pro	Phe	Arg	
		355						360					365				
5	aca	gaa	aag	gga	att	gtt	cga	aaa	tgg	atc	tca	aga	ttt	gaa	aaa	gtc	1152
	Thr	Glu	Lys	Gly	Ile	Val	Arg	Lys	Trp	Ile	Ser	Arg	Phe	Glu	Lys	Val	
		370					375					380					
10	tgg	cca	tac	ttg	gaa	acc	tac	aca	gag	gat	gtt	gct	cat	gaa	atc	tcc	1200
	Trp	Pro	Tyr	Leu	Glu	Thr	Tyr	Thr	Glu	Asp	Val	Ala	His	Glu	Ile	Ser	
		385				390					395				400		
15	aaa	gag	ttg	cac	ggc	acg	cca	gat	ctg	atc	atc	gga	aac	nac	agc	gac	1248
	Lys	Glu	Leu	His	Gly	Thr	Pro	Asp	Leu	Ile	Ile	Gly	Asn	Xaa	Ser	Asp	
					405					410					415		
	ggc	aat	atc	gtc	gcc	tcc	ttg	ctc	gca	cat	aaa	tta	ggg	gtc	aca	cag	1296
	Gly	Asn	Ile	Val	Ala	Ser	Leu	Leu	Ala	His	Lys	Leu	Gly	Val	Thr	Gln	
				420					425					430			
20	tgc	acc	atc	gcc	cat	gct	ttg	gag	aag	aca	aaa	tat	cca	gat	tca	gat	1344
	Cys	Thr	Ile	Ala	His	Ala	Leu	Glu	Lys	Thr	Lys	Tyr	Pro	Asp	Ser	Asp	
				435				440					445				
25	atc	tat	tgg	aag	aag	ctt	gaa	gac	aaa	tac	cat	ttc	tct	tgc	caa	ttt	1392
	Ile	Tyr	Trp	Lys	Lys	Leu	Glu	Asp	Lys	Tyr	His	Phe	Ser	Cys	Gln	Phe	
		450					455					460					
30	aca	gct	gat	ctt	ttt	gca	atg	aac	cat	aca	gat	ttc	atc	atc	acc	agt	1440
	Thr	Ala	Asp	Leu	Phe	Ala	Met	Asn	His	Thr	Asp	Phe	Ile	Ile	Thr	Ser	
		465				470					475				480		
35	act	ttc	cag	gaa	att	gca	gga	agc	aag	gac	act	gtt	ggg	caa	tac	gag	1488
	Thr	Phe	Gln	Glu	Ile	Ala	Gly	Ser	Lys	Asp	Thr	Val	Gly	Gln	Tyr	Glu	
					485					490					495		
	agc	cac	act	gct	ttc	act	ctt	cct	ggg	ctc	tac	cgt	gtt	gta	cat	ggg	1536
	Ser	His	Thr		500				Gly	Leu	Tyr	Arg	Val	Val	His	Gly	
								505						510			
40	atc	gat	gtg	ttt	gat	ccc	aaa	ttc	aac	att	gtt	tcc	cct	ggg	gct	gat	1584
	Ile	Asp	Val	Phe	Asp	Pro	Lys	Phe	Asn	Ile	Val	Ser	Pro	Gly	Ala	Asp	
			515					520					525				
45	atg	gag	ata	tac	ttc	cct	tac	acc	gaa	gag	aag	cgg	agg	ttg	aag	cat	1632
	Met	Glu	Ile	Tyr	Phe	Pro	Tyr	Thr	Glu	Glu	Lys	Arg	Arg	Leu	Lys	His	
			530				535					540					
50	ttc	cat	cct	gag	atc	gaa	gac	ctt	ctt	tac	acc	aaa	gtt	gag	aat	gaa	1680
	Phe	His	Pro	Glu	Ile	Glu	Asp	Leu	Leu	Tyr	Thr	Lys	Val	Glu	Asn	Glu	
		545				550					555					560	
55	gaa	cac	tta	tgt	gtg	ctc	aat	gac	cgc	aac	aag	cca	att	ctg	ttc	aca	1728
	Glu	His	Leu	Cys	Val	Leu	Asn	Asp	Arg	Asn	Lys	Pro	Ile	Leu	Phe	Thr	
					565				570						575		
	atg	cca	agg	ctt	gat	cgt	gtc	aag	aac	tta	acc	gga	ctc	gtc	gag	tgg	1776
	Met	Pro	Arg	Leu	Asp	Arg	Val	Lys	Asn	Leu	Thr	Gly	Leu	Val	Glu	Trp	
				580				585						590			
60	tgc	ggc	aag	aac	cca	aag	ttg	cgt	gag	ttg	gct	aac	ctc	gta	gtt	gta	1824
	Cys	Gly	Lys	Asn	Pro	Lys	Leu	Arg	Glu	Leu	Ala	Asn	Leu	Val	Val	Val	
			595					600					605				
	ggg	ggg	gat	agg	cga	aag	gaa	tct	aaa	gat	ttg	gaa	gag	aag	gct	gaa	1872

	Gly	Gly	Asp	Arg	Arg	Lys	Glu	Ser	Lys	Asp	Leu	Glu	Lys	Ala	Glu		
	610						615					620					
5	atg	aag	aaa	atg	ttt	gag	ctg	atc	gac	aag	tac	aac	ttg	aac	ggc	caa	1920
	Met	Lys	Lys	Met	Phe	Glu	Leu	Ile	Asp	Lys	Tyr	Asn	Leu	Asn	Gly	Gln	
	625					630					635				640		
10	ttc	aga	tgg	ata	tca	tct	caa	atg	aac	aga	atc	cga	aat	gtt	gaa	ctt	1968
	Phe	Arg	Trp	Ile	Ser	Ser	Gln	Met	Asn	Arg	Ile	Arg	Asn	Val	Glu	Leu	
					645					650					655		
15	tac	cga	tac	att	tgc	gac	acg	aaa	ggg	gcc	ttt	gta	cag	cct	gca	ttg	2016
	Tyr	Arg	Tyr	Ile	Cys	Asp	Thr	Lys	Gly	Ala	Phe	Val	Gln	Pro	Ala	Leu	
				660					665					670			
20	tat	gaa	gcc	ttt	gga	ttg	aca	gtt	gtg	gag	gca	atg	act	tgc	ggg	ttg	2064
	Tyr	Glu	Ala	Phe	Gly	Leu	Thr	Val	Val	Glu	Ala	Met	Thr	Cys	Gly	Leu	
			675					680					685				
25	cca	aca	ttc	gca	acc	tgt	aac	ggg	gga	cca	gcc	gag	att	att	gtc	cat	2112
	Pro	Thr	Phe	Ala	Thr	Cys	Asn	Gly	Gly	Pro	Ala	Glu	Ile	Ile	Val	His	
			690				695					700					
30	ggg	aaa	tct	ggg	ttc	aac	att	gat	cct	tac	cat	ggg	gat	caa	gct	gct	2160
	Gly	Lys	Ser	Gly	Phe	Asn	Ile	Asp	Pro	Tyr	His	Gly	Asp	Gln	Ala	Ala	
	705					710					715				720		
35	gac	ata	ctc	gtc	gat	ttc	ttt	gaa	aag	tgt	aag	aaa	gat	cca	tct	cac	2208
	Asp	Ile	Leu	Val	Asp	Phe	Phe	Glu	Lys	Cys	Lys	Lys	Asp	Pro	Ser	His	
					725					730					735		
40	tgg	gat	aag	atc	tcc	caa	gga	ggc	ttg	aaa	cga	ata	gag	gag	aag	tat	2256
	Trp	Asp	Lys	Ile	Ser	Gln	Gly	Gly	Leu	Lys	Arg	Ile	Glu	Glu	Lys	Tyr	
				740					745					750			
45	aca	tgg	aag	att	tac	tcg	gag	aga	cta	ttg	acc	ctg	aca	gga	gtg	tat	2304
	Thr	Trp	Lys	Ile	Tyr	Ser	Glu	Arg	Leu	Leu	Thr	Leu	Thr	Gly	Val	Tyr	
				755				760					765				
50	gga	ttc	tgg	aag	cat	gtt	tcc	aac	ctt	gaa	cgc	cgt	gag	agt	cgt	cgt	2352
	Gly	Phe	Trp	Lys	His	Val	Ser	Asn	Leu	Glu	Arg	Arg	Glu	Ser	Arg	Arg	
				770			775					780					
55	tac	ctt	gag														

5

<221> misc_feature
 <222> (414)..(414)
 <223> Xaa is Asn, Asp, His, or Tyr

5

<400> 2

Met ala Glu Arg Ala Leu Thr Arg Val His Ser Leu Arg Glu Arg Leu
 1 5 10 15

Asp Glu Thr Leu Leu Ala His Arg Asn Glu Ile Leu Ala Leu Leu Ser
 20 25 30

Arg Ile Glu Gly Lys Gly Lys Gly Ile Leu Gln His His Gln Ile Ile
 35 40 45

Leu Glu Phe Glu Ala Ile Pro Glu Glu Asn Arg Lys Lys Leu Ala Asn
 50 55 60

Gly Ala Phe Phe Glu Val Leu Lys Ala Ser Gln Glu Ala Ile Val Leu
 65 70 75 80

Pro Pro Trp Val Ala Leu Ala Val Arg Pro Arg Pro Gly Val Trp Glu
 85 90 95

Tyr Ile Arg Val Asn Val His Ala Leu Val Val Glu Glu Leu Thr Val
 100 105 110

Ala Glu Tyr Leu His Phe Lys Glu Glu Leu Val Asp Gly Ser Ser Asn
 115 120 125

Gly Asn Phe Val Leu Glu Leu Asp Phe Glu Pro Phe Asn Ser Ser Phe
 130 135 140

Pro Arg Pro Thr Leu Ser Lys Ser Ile Gly Asn Gly Val Glu Phe Leu
 145 150 155 160

Asn Arg His Leu Ser Ala Lys Leu Phe His Asp Lys Glu Ser Met His
 165 170 175

Pro Leu Leu Glu Phe Leu Arg Val His Cys His Lys Gly Lys Asn Met
 180 185 190

Met Leu Asn Asp Arg Ile Gln Asn Leu Asn Ala Leu Gln His Val Leu
 195 200 205

Arg Lys Ala Glu Glu Tyr Leu Gly Thr Leu Pro Pro Glu Thr Pro Cys
 210 215 220

Ala Glu Phe Glu His Arg Phe Gln Glu Ile Gly Leu Glu Arg Gly Trp

	225		230		235		240
5	Gly Asp Thr Ala	Glu Arg Val	Leu Glu Met	Ile Gln Leu	Leu Leu Asp		
		245	250		255		
10	Leu Leu Glu Ala	Thr Asp Pro Cys	Thr Leu Glu Lys	Phe Leu Gly Arg			
		260	265		270		
15	Ile Pro Met Val	Phe Asn Val	Val Ile Leu Thr	Pro His Gly Tyr	Phe		
		275	280		285		
20	Ala Gln Asp Asn	Val Leu Gly Tyr	Pro Asp Thr Gly	Gly Gln Val Val			
		290	295	300			
25	Tyr Ile Leu Asp	Gln Val Arg Ala	Leu Glu Asn Glu	Met Leu Leu Arg			
		305	310	315	320		
30	Ile Lys Gln Gln	Gly Leu Asn Ile	Thr Pro Arg Ile	Leu Ile Ile Thr			
		325	330	335			
35	Arg Leu Leu Pro	Asp Ala Val Gly	Thr Thr Cys Gly	Gln Arg Leu Glu			
		340	345	350			
40	Lys Val Tyr Gly	Thr Glu His Ser	Asp Ile Leu Arg	Val Pro Phe Arg			
		355	360	365			
45	Thr Glu Lys Gly	Ile Val Arg Lys	Trp Ile Ser Arg	Phe Glu Lys Val			
		370	375	380			
50	Trp Pro Tyr Leu	Glu Thr Tyr Thr	Glu Asp Val Ala	His Glu Ile Ser			
		385	390	395	400		
55	Lys Glu Leu His	Gly Thr Pro Asp	Leu Ile Ile Gly	Asn Xaa Ser Asp			
		405	410	415			
60	Gly Asn Ile Val	Ala Ser Leu Leu	Ala His Lys Leu	Gly Val Thr Gln			
		420	425	430			
65	Cys Thr Ile Ala	His Ala Leu Glu	Lys Thr Lys Tyr	Pro Asp Ser Asp			
		435	440	445			
70	Ile Tyr Trp Lys	Lys Leu Glu Asp	Lys Tyr His Phe	Ser Cys Gln Phe			
		450	455	460			
75	Thr Ala Asp Leu	Phe Ala Met Asn	His Thr Asp Phe	Ile Ile Thr Ser			
		465	470	475	480		
80	Thr Phe Gln Glu	Ile Ala Gly Ser	Lys Asp Thr Val	Gly Gln Tyr Glu			

	485	490	495
5	Ser His Thr Ala Phe Thr Leu Pro Gly Leu Tyr Arg Val Val His Gly 500 505 510		
10	Ile Asp Val Phe Asp Pro Lys Phe Asn Ile Val Ser Pro Gly Ala Asp 515 520 525		
15	Met Glu Ile Tyr Phe Pro Tyr Thr Glu Glu Lys Arg Arg Leu Lys His 530 535 540		
20	Phe His Pro Glu Ile Glu Asp Leu Leu Tyr Thr Lys Val Glu Asn Glu 545 550 555 560		
25	Glu His Leu Cys Val Leu Asn Asp Arg Asn Lys Pro Ile Leu Phe Thr 565 570 575		
30	Met Pro Arg Leu Asp Arg Val Lys Asn Leu Thr Gly Leu Val Glu Trp 580 585 590		
35	Cys Gly Lys Asn Pro Lys Leu Arg Glu Leu Ala Asn Leu Val Val Val 595 600 605		
40	Gly Gly Asp Arg Arg Lys Glu Ser Lys Asp Leu Glu Glu Lys Ala Glu 610 615 620		
45	Met Lys Lys Met Phe Glu Leu Ile Asp Lys Tyr Asn Leu Asn Gly Gln 625 630 635 640		
50	Phe Arg Trp Ile Ser Ser Gln Met Asn Arg Ile Arg Asn Val Glu Leu 645 650 655		
55	Tyr Arg Tyr Ile Cys Asp Thr Lys Gly Ala Phe Val Gln Pro Ala Leu 660 665 670		
60	Tyr Glu Ala Phe Gly Leu Thr Val Val Glu Ala Met Thr Cys Gly Leu 675 680 685		
65	Pro Thr Phe Ala Thr Cys Asn Gly Gly Pro Ala Glu Ile Ile Val His 690 695 700		
70	Gly Lys Ser Gly Phe Asn Ile Asp Pro Tyr His Gly Asp Gln Ala Ala 705 710 715 720		
75	Asp Ile Leu Val Asp Phe Phe Glu Lys Cys Lys Lys Asp Pro Ser His 725 730 735		
80	Trp Asp Lys Ile Ser Gln Gly Gly Leu Lys Arg Ile Glu Glu Lys Tyr		

	740	745	750
5	Thr Trp Lys Ile Tyr Ser Glu Arg Leu Leu Thr Leu Thr Gly Val Tyr 755 760 765		
10	Gly Phe Trp Lys His Val Ser Asn Leu Glu Arg Arg Glu Ser Arg Arg 770 775 780		
	Tyr Leu Glu Met Phe Tyr Ala Leu Lys Tyr Arg Lys Leu Ala Glu Ser 785 790 795 800		
15	Val Pro Leu Ala Glu Glu 805		

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU01/01580

A. CLASSIFICATION OF SUBJECT MATTER

Int. Cl. ⁷: A01H 5/00 C12N 15/11 C12N 15/29 C12N 15/54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

WPIDS, CA

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SEE ELECTRONIC DATABASES

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPIDS (A01H + keywords); CA, AGRICOLA, MEDLINE (sucrose synthase, susy, transgen?, transform?, recombinant, plant, cotton, fibre, fiber, seed)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Ruan, Y-L. <i>et al.</i> 2000. Pathway and control of sucrose import into initiating cotton fiber cells. Australian Journal of Plant Physiology. 27:795-800. See whole document, in particular "Results and discussion" pp797-800.	1-30
X	Ruan, Y-L. and Chourey, P.S. 1998. A fiberless seed mutation in cotton is associated with lack of fiber cell initiation in ovule epidermis and alterations in sucrose synthase expression and carbon partitioning in developing seeds. Plant Physiology. 118:399-406. See whole document, in particular "Results" and "Discussion" pp400-405.	1-13, 17-30
A	Chengappa, S. <i>et al.</i> 1999. Transgenic tomato plants with decreased sucrose synthase are unaltered in starch and sugar accumulation in the fruit. Plant Molecular Biology. 40:213-221. See "Abstract" and "Results and Discussion".	1-13, 17-30

☒ Further documents are listed in the continuation of Box C ☐ See patent family annex

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search
15 January 2002

Date of mailing of the international search report 31 JAN 2002

Name and mailing address of the ISA/AU
AUSTRALIAN PATENT OFFICE
PO BOX 200, WODEN ACT 2606, AUSTRALIA
E-mail address: pct@ipaustalia.gov.au
Facsimile No. (02) 6285 3929

Authorized officer

Terry Moore
Telephone No : (02) 6283 2632

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU01/01580

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	D'Aoust, M-A. <i>et al.</i> 1999. Antisense inhibition of tomato fruit sucrose synthase decreases fruit setting and the sucrose unloading capacity of young fruit. <i>The Plant Cell</i> . 11:2407-2418. See "Abstract" and "Discussion".	1-13, 17-30
A	Ruan, Y-L. <i>et al.</i> 1997. The differential expression of sucrose synthase in relation to diverse patterns of carbon partitioning in developing cotton seed. <i>Plant Physiology</i> . 115:375-385. See "Introduction" and "Discussion".	1-30
A	Shimizu, Y. <i>et al.</i> 1997. Changes in levels of mRNAs for cell wall-related enzymes in growing cotton fiber cells. <i>Plant and Cell Physiology</i> . 38(3):375-378. See whole document.	1-30
T	Haigler, C.H. <i>et al.</i> 2001. Carbon partitioning to cellulose synthesis. <i>Plant Molecular Biology</i> . 47:29-51. See whole document.	1-13, 17-30